

Detection of Genotoxicity of Phenolic Antioxidants, Butylated hydroxyanisole and *tert*-butylhydroquinone in Multiple Mouse Organs by the Alkaline Comet Assay

Ramadan, A.M. Ali^{1,2,*}, Takayoshi Suzuki²

¹Zoology Dept., College for Girls for Science, Arts and Education, Ain-Shams Univ., Heliopolis, Cairo, Egypt

²Division of Genetics and Mutagenesis, National Institute of Health Sciences, 1-18-1, Kamiyoga, Setagaya-ku, Tokyo 158, Japan

ramadanali27@gmail.com

Abstract: In this study we tested the genotoxicity of three widely used phenolic antioxidants, butylated hydroxyanisole (BHA) and *tert*-butylhydroquinone (t-BHQ) in multiple mouse organs using the alkaline comet assay. Tissue samples from four organs (stomach, liver, kidney and bone marrow) were collected from mice at 3 and 24 hrs post treatment with BHA (800 mg/kg) or t-BHQ (400 mg/kg) and examined for genotoxicity. The two compounds induced significant increase in DNA migration in a time dependant manner in specific organs. Extensive DNA damage was observed in stomach cells at 24 hrs post treatment in treatment groups. In addition to stomach, t-BHQ treatment induced significant increase in DNA migration in liver and kidney cells also. Although increased DNA damage was found in kidney cells of treatment groups at 3 hrs time point, at later time point it was persistent only in mice treated with tBHQ and in other treatment group (BHA) it appeared to be recovered with time. Evidently, bone marrow cells did not show genotoxicity in response to treatment with t-BHQ and BHA.

[Ramadan, A.M. Ali and Takayoshi Suzuki **Detection of Genotoxicity of Phenolic Antioxidants, Butylated hydroxyanisole and *tert*-butylhydroquinone in Multiple Mouse Organs by the Alkaline Comet Assay**] Life Science Journal 2012;9(1):177-183]. (ISSN: 1097-8135). <http://www.lifesciencesite.com>. 25

Key words: BHA; t-BHQ; comet assay; mice; bone marrow; liver; kidney; stomach; ENU.

1. Introduction

Butylated hydroxyanisole (BHA, 3-*tert*-butyl 4-hydroxyanisole) and its *O*-demethylated metabolite *tert*-butylhydroquinone (t-BHQ) are synthetic phenolic food antioxidants widely used to protect oils, fats and shortenings from oxidative deterioration and rancidity (Aluyor Ori-Jesu, 2008; Finley *et al.*, 2011). Both compounds are generally non-mutagenic and nearly all short-term genotoxicity tests including the classic *Salmonella typhimurium* (with or without S9 fraction), unscheduled DNA synthesis assay using rat hepatocytes, and except both positive and negative results with and without activation respectively in Chinese hamster cells for chromosomal aberration and sister chromatid exchange are negative (Rogers *et al.*, 1985; Williams *et al.*, 1990). However, chronic dietary administration of BHA resulted in papilloma and carcinoma formation in the forestomachs of rats, mice and hamsters (IARC, 1986; Clayson *et al.*, 1990; Chandra *et al.*, 2010). In fact, both BHA and t-BHQ have shown to contain both carcinogenic and anticarcinogenic properties (Whysner *et al.*, 1994; Hirose *et al.*, 1997; Kahl, 1997; Gharaviet *et al.*, 2007). While, anticarcinogenic activity of BHA has been ascribed to its ability to induce Phase II metabolic enzymes and its free radical trapping activity (Williams and Iatropoulos, 1997; Li *et al.*, 2005; Kadoma *et al.*, 2008; Cemeli and Anderson, 2011), its carcinogenicity has been often linked to its

quinone-forming metabolites, a property that BHA shares with benzene. Quinones are highly reactive molecules, which can bind to biological macromolecules (O'Brien, 1991; Tuet *et al.*, 2011) and induce chromosomal loss or chromosomal breakage (Dobo *et al.*, 1994; Jacobus *et al.*, 2008).

The scope of this research is to determine the *in vivo* genotoxic effects of t-BHQ and BHA in different organs of mouse. In this study we conducted the single cell gel electrophoresis assay (Comet assay) on four organs (stomach, liver, kidney and bone marrow) of mice treated with t-BHQ (400 mg/kg) or BHA (800 mg/kg). To identify possible time dependent effect, tissue samples were collected on two occasions, 3 and 24 hrs post treatment and subjected to the alkaline version of the comet assay. The comet assay is a sensitive cytogenetic assay with advantages of detecting broad spectrum of DNA damages which may lead into DNA strand breaks, the need of small number of cells per sample, and the short time needed to complete a study (Tice *et al.*, 2000). In addition, the comet assay can be applied to any tissue in the given *in vivo* model, and has potential advantages over other *in vivo* genotoxicity tests methods, which are reliably applicable to one or few tissues. This is very important for investigation of suspected tissue specific-genotoxic activity, which include 'site of contact' genotoxicity.

Material and Methods

Chemicals

All chemicals used in this study were purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan).

Animals and treatment.

Male ddY mice were obtained from Japan SLC (Shizuoka, Japan) at 7 weeks of age and used after one week of acclimatization. They were maintained at 20–24°C and 55–65% humidity with a 12-hr light-dark cycle and fed commercial pellets (Oriental Yeast Industries Co., Tokyo, Japan) and tap water *ad libitum*. Aqueous solutions of t-BHQ (400 mg/kg) and BHA (800 mg/kg) were administered intragastrically to groups of three mice at a dose level of 10 ml/kg. A positive control (ENU, 20 mg/kg) group of two mice were treated at the same time in the same manner. The animals were sacrificed by cervical dislocation at 3 hr and 24 hrs after treatment and four organs - bone marrow, liver, kidney and stomach - were dissected out. Tissue samples were collected from four untreated mice at the latter sampling point (24 hrs) and treated as the negative control.

The tissue samples except bone marrow were washed in saline, minced, suspended at a concentration of 1 g/ml in ice cold homogenizing buffer (HBSS with 20 mM EDTA, 10% DMSO, pH 7.5) and homogenized gently using a Dounce's homogenizer. The marrow was collected from the femur bones and suspended in 1 ml of chilled homogenizing buffer. The cell suspensions were diluted in chilled homogenizing buffer appropriately and subjected to the alkaline comet assay immediately.

Comet assay

The alkaline comet assay was performed basically as described by Miyamae *et al.* (1998) with few modifications. The cell suspensions prepared as above were mixed 1:1 (v/v) with 1% low melting point (37°C) agarose (LMA) prepared in PBS, pH 7.4 and 75 µl of the mixture was quickly layered on a 1% normal melting point agarose (NMA) (prepared in distilled water) precoated and overnight dried slides and covered with a coverslip. Then the slides were placed on a chilled plate to allow complete polymerization of agarose. Finally 75 µl of 0.5% NMA in PBS was quickly layered in the same manner after removing the coverslip and allowed to solidify on chilled plate. The slides were immersed in freshly made chilled lysis solution (2.5M NaCl, 100mM Na₂EDTA, 10mM Trizma base, 10% DMSO and 1% Triton X-100; pH 10.0) in dark at 4°C for 60 min. The slides were then placed in a horizontal electrophoresis tank containing electrophoresis buffer (300mM NaOH and 1mM Na₂EDTA; pH 13.0) for

10 min, allowing salt equilibration and further DNA unwinding before electrophoresis at 25 V (0.8V/cm), 300 mA for 20 min. The slides were washed three times (5 min each) with chilled neutralizing buffer, 0.4M Tris-HCl (pH 7.5). After the third wash, the slides were stained with 50 µl EtBr (20 µl/ml) and covered with a coverslip. To prevent drying, the slides were stored in a humidified container until microscopic examination.

The slides were examined at 200x magnification using Olympus fluorescent microscope. All slides were coded and examined blindly. A total of 1000 randomly selected cells from two replicate slides (500 cells per slide) were examined per sample. The comets were classified into five categories (Type 1 – Type 5) depending on the fraction of DNA migrated out into the tail, and thus increasing degrees of damage (Miyamae *et al.*, 1998). The comets were assigned a value of 0, 1, 2, 3 or 4 (from undamaged, 0, to maximally damaged, 4). Thus, the total score for 1000 comets ranged from 0 (all undamaged) to 4000 (all damaged) in arbitrary units (Visvardis *et al.*, 1997; Piperakis *et al.*, 1998). In this method a large number of cells can be examined in a short time.

Statistical analysis

Results of the different treatment groups were compared to data obtained from four untreated mice using Students' one-tailed *t*-test. Significance was indicated by *P* values <0.05.

3. Results


Table 1 presents the comet data measured by visual scoring from four organs (bone marrow, liver, kidney and stomach) from mice intragastrically treated with t-BHQ (400mg/kg b.w.) or BHA (800 mg/kg b.w.) for 3 and 24 hr. In this experiment comet data obtained from four untreated mice were treated as the negative control and used for comparison. Among t-BHQ treated animals, 24 hr after exposure mean DNA migration score was significantly (*P*<0.05) high in 3 organs (stomach, liver and kidney), although at earlier time point relative to control value it was different significantly only in kidney. In bone marrow the induced DNA strand breaks were not significant at both time points.

Exposure to BHA resulted in significant increase in DNA migration in a time dependent manner in specific organs. Although significant (*P*<0.05) increase in DNA damage was observed among stomach tissues at both sampling points, in kidney it occurred only at 3 hr time point and then decreased with time. Following BHA treatment liver and bone marrow cells did not yield any significant increase in mean DNA migration. ENU (20 mg/kg) treatment

produced highly significant ($P < 0.001$) DNA damage in all four tested organs at both time points.

Figure 1 shows the distribution pattern of nuclear DNA expressed as the percent cells in the five comet classes from Type 1 (undamaged) to Type 5 (maximally damaged) in various organs of mice belonged to different treatment groups. Except in ENU (positive control) treated group, in other samples some cells/nuclei remained undamaged (category Type 1). At both sampling time points in all treatment groups greater numbers of comets showed high levels of DNA damage (category Type

4) in stomach cells. The frequency distribution of DNA damage among individual hepatocytes showed that the number of damaged cells and the extent of damage were greatest among mice treated with tBHQ (Figs. 1-C and 1-D). Similarly in renal cells the extent of DNA damage decreased with time in BHA treated animals (Figs. 1-E and 1-F). Higher levels of damage were observed in renal cells from t-BHQ treated mice. Among bone marrow cells in treatment groups (t-BHQ and BHA) over 50% of the cells showed no damage (Figs. 1-A and 1-B).



G

H




Fig. 1. Incidences of comet cells in mice treated with t-BHQ (400 mg/kg b.w.), BHA (800 mg/kg b.w.) and the positive control ENU (20 mg/kg b.w.) for 3 and 24 h. The comet cells derived from bone marrow (Figs. 1A and 1B), liver (Figs. 1C and 1D), kidney (Figs. 1E and 1F) and stomach (Figs. 1G

Table 1. Incidences of comet cells without type 1 in bone marrow, liver, kidney and stomach of mice treated with t-BHQ and BHA for 3 and 24 h.

Treatment	Pooled calculations of incidences of comets without type 0 (Average/500 cells \pm S.D.)							
	Bone marrow		Liver		Kidney		Stomach	
	3 h	24 h	3 h	24 h	3 h	24 h	2h	24 h
Control (Dis. H ₂ O)	146.3 \pm 125.3	156.5 \pm 16.3	202.0 \pm 116.0	369.5 \pm 99.3	283.3 \pm 110.3	305.3 \pm 94.6	350.0 \pm 109.4	479.5 \pm 14.5
t-BHQ (400 mg/kg)	213.7 \pm 136	234.3 \pm 37.7	274.5 \pm 61.3	425.2 \pm 59.6*	447.5 \pm 66.1	385.2 \pm 66.1*	453.5 \pm 29.1	490.3 \pm 8.1*
BHA (800 mg/kg)	279.3 \pm 171.5	226.0 \pm 45.1	345.0 \pm 40.5	237.2 \pm 72.2	397.2 \pm 79.1*	287.3 \pm 97.7	483.3 \pm 13.6*	490.1 \pm 11.6*
ENU (20 mg/kg)	495.0 \pm 6.0**	498.5 \pm 3.0**	500.0 \pm 0.0**	500.0 \pm 0.0**	500.0 \pm 0.0**	500.0 \pm 0.0**	498.25 \pm 2.1**	500.0 \pm 0.0**

n.b.

ns = non significant; * = (P<0.05); * = (P<0.01); ** = (P<0.001)

4. Discussion

Considering the strong correlation between organ specific genotoxicity and organ specific carcinogenicity, the assessment of genotoxicity in multiple organs *in vivo* may indicate would be target organ(s) in humans and provide useful information for the evaluation of chemical safety. While other *in vivo* genotoxicity tests are limited to one or few tissues, the comet assay can be applied to any tissue provided that a single cell/nucleus suspension can be obtained. Therefore, in the present research, to test the tissue specific genotoxic effect of t-BHQ and, BHA mice were administered intragastrically with the test compounds and various organs were examined for genotoxicity using the comet assay. Results of this research suggest that the test compounds induced significant DNA strand breaks in stomach cells. The effect was more pronounced among BHA treated animals at both sampling occasions. With t-BHQ, a significantly higher degree of DNA damage was observed at 24 hr time point, although at earlier time point (3 hrs) the induced DNA damage in these treatment groups were not significantly different from control data. These observations again acknowledge the earlier findings

that stomach in particular forestomach in rodent is the target organ for BHA induced genotoxicity. It is also important to remember that there are several bioassay reports describing neoplasm-promoting effects of BHA and t-BHQ in the fore stomach when given after initiating carcinogens such as N-methyl-N-nitrosoguanidine (MNNG) (Takahashi *et al.*, 1986, Whysner *et al.*, 1994; Miyauchi *et al.*, 2002; Kuroiwa *et al.*, 2007; Liu and Russell, 2008), N-methyl-N-nitrosourea (NMU) (Imaida *et al.*, 1984; Tsuda *et al.*, 1984; García-González *et al.*, 2000), N,N-dibutyl nitrosamine (DBN) (Fukushima *et al.*, 1987). In this respect, there are studies, which demonstrated that BHA inhibits cell-to-cell communication (Williams *et al.*, 1990), which is often considered a marker for tumor promoters (Budunova and Williams, 1994; Yamasaki, 1996; Leithe *et al.*, 2006; Vinken *et al.*, 2009). In the bone marrow cells, although at earlier time point a non-significant increase in DNA migration was observed in BHA and t-BHQ treated animals, it was not persistent at latter time point and appeared to be recovered with time. *In vivo* BHA is metabolized to t-BHQ in the liver (Peters *et al.*, 1996; Moridani *et al.*, 2002). t-BHQ and HQ are further oxidized to

their respective quinone species, 1,4-benzoquinone and tertiary-butylquinone, which in turn may enter into a redox cycling between quinone and hydroquinone via semiquinone radicals and can generate reactive oxygen species (Peters *et al.*, 1996; Li *et al.*, 2002). These ROS may then react with cellular protein and DNA and thus cause large-scale chromosomal alterations (Cerutti, 1985). The test compounds used in this study have been reported to be nephrotoxic and can induce tumor in rodent kidney (Kari *et al.*, 1992; Shibata *et al.*, 1991; Tsuda *et al.*, 1984). Nonetheless, the evidence for genotoxicity as increased level of DNA migration in the liver and kidney was observed only in mice treated with t-BHQ. The nephrotoxicity of these compounds are often linked to their quinonethioether metabolites (Monks *et al.*, 1992) and their ability to react with DNA either directly (Jacobus *et al.*, 2008) or indirectly via reactive oxygen generation (Canales *et al.*, 1993; Li *et al.*, 2002; Li *et al.*, 2007). Hence, differences in metabolism of these compounds and ability of their metabolites to generate ROS could be a possible reason for the observed differences in toxicity (Kadoma *et al.*, 2008).

DNA strand breaks are associated both to the removal of DNA adducts by endonuclease and to the direct action of the chemicals and/or the free radicals produced during their metabolism, which then creates both single- and double-strand breaks in DNA (Bohr, 1991). Investigations of BHA and its metabolites have not demonstrated DNA adduct formation, as measured by the sensitive [³²P]postlabelling assay (Saito *et al.*, 1989). Previous research has showed that enzymatic oxidation of t-BTQ in rodent forestomach may occur via prostaglandin H synthase PHS (Schilderman *et al.*, 1993) or possibly a gastric peroxidase (Banerjee, 1988). It is widely believed that t-BHQ, a *O*-demethylated metabolite of BHA probably produces many of the toxic effects ascribed to BHA. It is also postulated that BHA and tBHQ act through ROS induced oxidative damage including DNA damage (Inverson, 1995). Supporting this suggestion studies have demonstrated the formation of 8-hydroxydeoxyguanosine, a marker for reactive oxygen formation and for DNA strand breaks (Okubo *et al.*, 1997; Schilderman *et al.*, 1995), in calf thymus DNA (Nagai *et al.*, 1996) and in isolated DNA and cultured rat hepatocytes (Li *et al.*, 2002) treated with tBHQ. Peters *et al.* (1996) have demonstrated that tBHQ and its glutathione conjugates are capable of catalyzing 8-hydroxydeoxyguanosin formation and thus induce oxidative DNA damage. Further, Dobo and Eastmond (1994) have demonstrated the role of reactive oxygen species in the chromosomal breakage induced by tBHQ in a prostaglandin H synthase containing Chinese hamster

V79 cell line. Considering these findings, although tBHQ and BHA are generally considered nongenotoxic, the DNA damage observed in this experiment may be related to their indirect action on DNA via ROS mechanism. Since toxicity of these compounds are often ascribed to their metabolic products such as quinonethioethers and hence differences in the metabolism of these compounds may play an important role in determining the target organ of toxicity.

Corresponding author

Ramadan A. M. Ali
Department of Zoology, Univ. College for Women,
Ain Shams University, Egypt
ramadanali27@gmail.com

References

- Aluyor E. and Ori-Jesu M. (2008). The use of antioxidants in vegetable oils – A review. *Afri. J. Biotechnol.*, 7 (25): 4836-4842.
- Banerjee K. (1988). Membrane peroxidase. *Mol. Cell Biochem.*, 83: 105-128.
- Bohr A. (1991). Gene specific DNA repair. *Carcinogenesis*, 12: 1983-1992.
- Budunova I.V. and Williams G.M. (1994). Cell culture assays for chemicals with tumor-promoting or tumor-inhibiting activity based on the modulation of intercellular communication. *Cell Biol. Toxicol.*, 10: 71-116.
- Canales P.L., Kleiner H.E., Monks T.J. and Lau S.S. (1993). Formation of 8-hydroxydeoxyguanosine by quinone-thioethers. *Toxicologists*, 13: 202 (Abstract).
- Cemeli E. and Anderson D. (2011). Mechanistic investigation of ROS-induced DNA damage by oestrogenic compounds in lymphocytes and sperm using the comet assay. *Int. J. Mol. Sci.*, 12: 2783-2796.
- Cerutti P.A. (1985). Prooxidant state and tumor promotion. *Science*, 227: 375-381.
- Chandra S., Nolan M. and Malarkey D. (2010). Chemical carcinogenesis of the gastrointestinal tract in rodents: An overview with emphasis on NTP carcinogenesis. *Bioassays Toxicol. Pathol.*, 38: 188-197.
- Clayson D.B., Iverson F., Nera E.A. and Lok E. (1990). The significance of induced forestomach tumors. *Ann. Rev. Pharmacol. Toxicol.*, 30: 441-463.
- Dobo K.L. and Eastmond D.A. (1994). Role of oxygen radicals in the chromosomal loss and breakage induced by the quinone-forming compounds, hydroquinone and *tert*-butylhydroquinone. *Environ. Mol. Mutagen.*, 24: 293-300.

- Finley J. Kong A., Hintze K., Jeffery E., Ji L. and Lei X. (2011). Antioxidants in foods: state of the science important to the food industry. *J. Agric. Food Chem.*, 59 (13): 6837–6846
- Fukushima S., Ogiso T., Kurata Y., Hirose M. and Ito N. (1987). Dose-dependent effects of butylatedhydroxytoluene and ethoxyquin for promotion of bladder carcinogenesis in N-butyl-N-(4-hydroxybutyl) nitrosamine-initiated, unilaterally ureter ligated rats. *Cancer Letters*, 34: 83-90.
- Gharavi N., Haggarty S. and El-Kadi A. (2007). Chemoprotective and carcinogenic effects of tert-butylhydroquinone and its metabolites. *Current Drug Metabolism*, 8: 1-7.
- Hirose M., Takesada Y., Tanaka H., Tamano S., Kato T. and Shirai T. (1997). Carcinogenicity of antioxidants BHA, caffeic acid, sesamol, 4-methoxyphenol and catechol at low doses, either alone or in combination and modulation of their effects in rat medium-term multiorgan carcinogenesis model. *Carcinogenesis*, 19: 207-212.
- IARC (1986). IARC monographs on the evaluation of the carcinogenic risks of chemicals to humans. vol. 40, some naturally occurring and synthetic food components, furocoumarins and ultraviolet radiation butylatedhydroxyanisole (bha), International Agency for Research on Cancer, Lyon. pp. 123-159.
- Iverson F. (1995) Phenolic antioxidants: health protection branch studies on butylatedhydroxyanisole. *Cancer Letters*, 93: 49-54.
- Jacobus J., Flor S., Klingelutz A., Robertson L., and Ludewig G. (2008). 2-(4'-chlorophenyl)-1,4-benzoquinone increases the frequency of micronuclei and shortens telomeres. *Environ Toxicol Pharmacol.*, 25(2): 267–272.
- Kadoma Y., Ito S., Yokoe I. and Fujisawa S. (2008). Comparative study of the alkyl and peroxy radical-scavenging activity of 2-*t*-butyl-4-methoxyphenol (bha) and its dimer, and their theoretical parameters. *In vivo*, 22: 289-296.
- Kahl R. (1997) Synthetic antioxidants: biochemical actions and interferences with radiation and toxic compounds, chemical mutagens and chemical carcinogens. *Toxicology*, 33: 185-228.
- Kari F.W., Bucher J., Eustis S.L., Haseman J.K. and Huff J.E. (1992). Toxicity and carcinogenicity of hydroquinone in F344/N rats and B6C3F₁ mice. *Food Chem. Toxicol.*, 30: 737-747.
- Kuroiwa, Y., Ishii, Y., Umemura, T., Kanki, K., Mitsumori, K., Nishikawa, A., Nakazawa, H. and Hirose, M. (2007). Combined treatment with green tea catechins and sodium nitrite selectively promotes rat forestomach carcinogenesis after initiation with *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine. *Cancer Science*, 98: 949–957.
- Leithe E., Sirnes S., Omori Y. and Rivedal E. (2006). Downregulation of gap junctions in cancer cells. *Critical Reviews TM in Oncogenesis*, 12(3-4): 225–256
- Li Y., Zhong F., Wu S. and Shi N. (2007). NF-E2 related factor 2 activation and heme oxygenase-1 induction by tert-butylhydroquinone protect against deltamethrin-mediated oxidative stress in PC12 cells. *Chem. Res. Toxicol.*, 20(9): 1242-51.
- Li Y., Seacat A., Kuppasamy P., Zweier J.L., Yager J.D. and Trush, M.A. (2002). Copper redox-dependent activation of 2-*tert*-butyl(1,4)hydroquinone: formation of reactive oxygen species and induction of oxidative DNA damage in isolated DNA and cultured rat hepatocytes. *Mutat. Res.*, 518: 123-133.
- Li J., Johnson D., Calkins M., Wright L., Svendsen C. and Johnson J. (2005). Stabilization of Nrf2 by *t*BHQ confers protection against oxidative stress-induced cell death in human neural stem cells. *Toxicol. Sci.*, 83: 313-328.
- Liu C. and Russell R. M. (2008). Nutrition and gastric cancer risk: an update. *Nutr. Rev.*, 66: 237–249.
- Miyamae Y., Yamamoto M., Sasaki Y.F., Kobayashi H., Igarashi-Soga M., Shimoi K. and Hayashi M. (1998). Evaluation of a tissue homogenization technique that isolates nuclei for the in vivo single cell gel electrophoresis (comet) assay: a collaborative study by five laboratories. *Mutat. Res.*, 418: 131-140.
- Monks T.J., Hanzlik R.P., Cohen G.M., Ross D. and Graham D.G. (1992). Quinone chemistry and toxicity. *Toxicol. App. Pharmacol.*, 112: 2-16.
- Moridani M., Cheon S., Khan S. and O'Brien J. (2002). Metabolic activation of 4-hydroxyanisole by isolated rat hepatocytes. *Drug Metab. Dispos.*, 30(10): 1063-9.
- Nagai F., Okubo T., Ushiyama K., Satoh K. and Kano I. (1996). Formation of 8-hydroxydeoxyguanosine in calf thymus DNA treated with *tert*-butylhydroquinone, a major metabolite of butylatedhydroxyanisole. *Toxicol. Lett.*, 89: 163-167.
- O'Brien P.J. (1991). Molecular mechanisms of quinone cytotoxicity. *Chem. Biol. Inter.*, 80: 1-41.
- Okubo T., Nagai F., Ushiyama K. and Kano I. (1997). Contribution of oxygen radicals to DNA cleavage by quinone compounds derived from phenolic antioxidants, *tert*-butylhydroquinone and 2,5-di-*tert*-butylhydroquinone. *Toxicol. Lett.*, 90: 11-18.
- Peters M., Lau S., Dulik D., Murphy D., Ommen B., Bladeren P. and Monks T. (1996). Metabolism of *tert*-Butylhydroquinone to *S*-Substituted Conjugates

- in the Male Fischer 344 Rat. *Chem. Res. Toxicol.*, 9: 133-139.
- Piperakis S.M., Visvardis E-E., Sagnou M. and Tassiou A.M. (1998). Effects of smoking and ageing on oxidative DNA damage of human lymphocytes. *Carcinogenesis*, 19: 695-698.
- Rogers C.G., Nayak B.N. and Heroux-Metcalf C. (1985). Lack of induction of sister chromatid exchanges and of mutation to 6-thioguanine resistance in V79 cells by butylatedhydroxyanisole with and without activation by rat or hamster hepatocytes. *Cancer Lett.*, 27: 61-69.
- Saito K., Nakagawa S., Yoshitake A., Miyamoto J., Hirose M. and Ito N. (1989). DNA-adduct formation in the forestomach of rats treated with 3-tert-butyl-4-hydroxyanisole and its metabolites as assessed by an enzymatic ³²P-postlabeling method. *Cancer Lett.*, 48: 189-195.
- Schilderman P.A.E.L., Rhijnsburger E., Zwingmann I. and Kleinjans J.C.S. (1995). Induction of oxidative DNA damage and enhancement of cell proliferation in human lymphocytes *in vitro* by butylatedhydroxyanisole. *Carcinogenesis* 16: 507-512.
- Schilderman P.A.E.L., van Maanen J.M.S., Smeets E.J., ten Hoor F. and Kleinjans J.C.S. (1993). Oxygen radical formation during prostaglandin H synthase-mediated biotransformation of butylatedhydroxyanisole. *Carcinogenesis*, 14: 347-353.
- Shibata M.A., Asakawa E., Hagiwara A., Kurata Y. and Fukushima S. (1991). DNA synthesis and scanning electron microscopic lesions in renal pelvic epithelium of rats treated with bladder cancer promoters. *Toxicol. Lett.*, 55: 263-272.
- Takahashi M., Furukawa F., Toyoda K., Sato H., Hasegawa R. and Hayashi Y. (1986). Effects of four antioxidants on N-methyl-N'-nitro-N-nitrosoguanidine initiated gastric tumor development in rats. *Cancer Lett.*, 30: 161-168.
- Tice R.R., Agurell E., Anderson D., Burlinson B., Hartmann A., Kobayashi H., Miyamae Y., Rojas E., Ryu J.C. and Sasaki Y.F. (2000). The single cell gel/comet assay: guidelines for *in vitro* and *in vivo* genetic toxicology testing. *Environ. Mol. Mutagen.*, 35: 206-221.
- Tsuda H., Fukushima S., Imaida K., Sakata T. and Ito N. (1984). Modification of carcinogenesis by antioxidants and other compounds. *Acta Pharmacol. Toxicol.*, 2(Suppl.), 125-143.
- Tu T., Giblin D. and Gross M. (2011). Structural Determinant of Chemical Reactivity and Potential Health Effects of Quinones from Natural Products. *Chem. Res. Toxicol.*, 24 (9): 1527-1539
- Vinken M., Doktorova T., Decroock E., Leybaert L., Vanhaecke T. and Rogiers V. (2009). Gap junctional intercellular communication as a target for liver toxicity and carcinogenicity. *Crit. Rev. Biochem. Mol. Biol.*, 44(4):201-222.
- Visvardis E-E., Tassiou A.M. and Piperakis S.M. (1997). Study of DNA damage induction and repair capacity of fresh and cryopreserved lymphocytes exposed to H₂O₂ and γ -radiation with the alkaline comet assay. *Mutat. Res.*, 383: 71-80.
- Whysner J., Wang C., Zang E., Latropoulos M.J. and Williams G.M. (1994). Dose-response promotion by butylatedhydroxyanisole in chemically initiated tumors of the rodent forestomach. *Food Chem. Toxicol.*, 32: 215-222.
- Williams G. M. and Iatropoulos (1997). Anticarcinogenic effects of synthetic phenolic antioxidants. In *Oxidants, Antioxidants, and Free Radicals*, ed. S.I. Baskin and H. Salem, pp. 341-350. Taylor & Francis, New York.
- Williams G.M., McQueen C.A. and Tong C. (1990). Toxicity studies of butylatedhydroxyanisole and butylatedhydroxytoluene. I. Genetic and cellular effects. *Food Chem. Toxicol.*, 28: 793-798.
- Yamasaki H. (1996). Role of disrupted gap junctional intercellular communication in detection and characterization of carcinogens. *Mutat. Res.*, 365: 91-105.

12/20/2012