

## Impact of stone quarrying on the health of Residents in Nigeria

Igbinovia Osayi Martins, Osu Charles I \* and Iwuoha G.N.

\* Department of Pure and Industrial Chemistry, University of Port Harcourt, P.M.B 5323 Choba, Port Harcourt, Rivers State, Nigeria.

Telephone: +234 803 7783246 email: [charsike@yahoo.com](mailto:charsike@yahoo.com)

**Abstract:** This study investigated the effect of stone quarry activities and seasonal variation on the concentration of air pollutants. Higher concentration of particulate matter was observed in the dry season than wet season. Concentration of particulate matter ranged from  $10.40 \pm 0.02$  to  $18.70 \pm 1.40$  mg/m<sup>3</sup>, dry season;  $1.07 \pm 0.02$  to  $2.50 \pm 0.09$  mg/m<sup>3</sup>, wet season (Table 1) for site A and  $12.01 \pm 1.10$  mg/m<sup>3</sup> to  $13.60 \pm 0.30$  mg/m<sup>3</sup>, wet season;  $16.40 \pm 1.30$  to  $21.30 \pm 1.60$  mg/m<sup>3</sup>, dry season for site B. concentration of H<sub>2</sub>S ranges from  $0.30 \pm 0.05$  to  $1.70 \pm 0.10$  ppm, wet season;  $2.06 \pm 0.02$  to  $2.10 \pm 0.01$  ppm, dry season for site A and  $0.60 \pm 0.05$  to  $1.00 \pm 0.10$  ppm, wet season;  $3.17 \pm 0.02$  to  $1.30 \pm 0.01$  ppm, dry season for site B. Mean concentration of NO<sub>2</sub> ranges from  $0.50 \pm 0.01$  to  $1.90 \pm 0.03$  ppm, wet season;  $4.30 \pm 0.01$  to  $7.00 \pm 0.11$  ppm, dry season for site A and  $2.01 \pm 0.02$  to  $3.30 \pm 0.02$  ppm, wet season;  $4.70 \pm 0.01$  to  $6.30 \pm 0.10$  ppm, dry season for site B. Mean concentration of SO<sub>2</sub> in study locations shows the same trend of being higher during dry season than during wet season. This is due to dilution effect of rain on the atmosphere. Its values ranges from  $0.70 \pm 0.06$  to  $1.20 \pm 0.02$  ppm, wet season;  $5.14 \pm 0.20$  to  $6.10 \pm 0.08$  ppm, dry season for site A and  $1.30 \pm 0.02$  to  $5.10 \pm 0.15$  ppm, wet season  $6.20 \pm 0.10$  to  $6.80 \pm 0.04$  ppm, dry season for site B. The concentration of vanadium in the environment ranged from  $0.13 \pm 0.01$  to  $0.18 \pm 0.03$  and  $0.10 \pm 0.01$  to  $0.46 \pm 0.01$  for sample A and B respectively. The concentration of the heavy metal ranged from 14.00 to 42.50 mg/kg, Cu; 23.50 to 38.70 mg/kg, Cr; 45.60 to 136.30 mg/kg, Zn; 0.10 to 0.65 mg/kg, Hg; 10.60 16.80 mg/kg, Pb and 1.60 to 16.40 mg/kg, Ni for site A. Concentration of the heavy metal in site B ranged from 11.30 to 25.10 mg/kg, Cu; 4.70 to 15.20 mg/kg, Cr; 44.30 to 186.10 mg/kg, Zn; 0.01 to 0.20 mg/kg, Hg; 6.60 to 13.90 mg/kg, Pb and 1.00 to 5.40 mg/kg, Ni. This study found that about 99.99 % of air pollution diseases affect the households. Data from this study showed that chronic exposure to dust and polluted air from crushing of stones increased susceptibility to respiratory problems. Percentage values obtained were 24.49 %, catarrh; 28.57 %, coughing; 26.53 %, headache; 9.18 %, chest pains and 11.22 %, eye infections.

[Igbinovia O.M., Osu Charles I and Iwuoha G.N. **Impact of stone quarrying on the health of Residents in Nigeria.** *Life Sci J* 2017;14(10):59-64]. ISSN: 1097-8135 (Print) / ISSN: 2372-613X (Online). <http://www.lifesciencesite.com>. 9. doi:[10.7537/marslsj141017.09](https://doi.org/10.7537/marslsj141017.09).

**Keywords:** stone quarry; air parameters; health effect;

### 1. Introduction

The monitoring of vanadium (v) oxide and other air pollutants in ambient air has received substantial attention over the past several years because they are among the major pollutants which significantly affect the chemistry of the atmosphere and human health (Wu et al., 2003).

Vanadium is a non-volatile metal, and atmospheric transport is via particulates (Fish and Komlenic, 1984). Vanadium (V) oxide (V<sub>2</sub>O<sub>5</sub>) is a poorly soluble oxide which, in water or body fluids, releases some vanadium ions which may speciate either in cationic (VO<sup>2+</sup>) or anionic (HVO<sub>4</sub><sup>2-</sup>) forms. The adsorption, distribution and elimination of vanadium (V) oxide and other vanadium compounds are similar. There are variations depending on the solubility of the administered (National Toxicology Program, 2002). It is a yellow-orange powder or dark gray flakes (Bauer et al., 2003); yellow to rust-brown orthorhombic crystals (Lide, 2008); slightly soluble in

water, soluble in concentrated acids and alkalis; insoluble in ethanol (O' Neil, 2001).

Vanadium (V) oxide is used as an oxidation catalyst in heterogeneous and homogeneous catalytic processes for production of sulfuric acid from sulfur dioxide, phthalic anhydride etc. (O' Neil, 2001; Bauer et al., 2003). It is also used as a corrosion inhibitor in industrial process, as a coating for welding electrodes. Vanadium is a trace metal that is found naturally both on soil and water (Nriagu, 1998). Under environmental conditions, vanadium may exist in oxidation states +3, +4, and +5. V<sup>3+</sup> and V<sup>4+</sup> act as cations but V<sup>5+</sup>, the most common form in the aquatic environment, reacts both as a cation and anionically as an analogue of phosphate. Mineral dissolution rapidly oxidizes V<sup>3+</sup> and V<sup>4+</sup> to the pentavalent state. Vanadium has been characterized as a constituent of several enzymes systems and complexes within living organisms. Adsorption through the skin is thought to be quite minimal due to its low lipid/water solubility.

In human, acute vanadium poisoning can manifest itself in a number of symptoms including eye irritation and tremors of the hand, a greenish coloration of the tongue has been observed in humans exposed to high concentration of vanadium (V) oxide and is probably due to the formation of trivalent and tetravalent vanadium complexes; obstructive lung disease (National Toxicology Program, 2002); inflammatory lesions of the respiratory tract (Ress et al., 2003); chronic inflammation of the bronchi.

Vanadium is a by-product of oil-burning furnaces when vanadium pentoxide ( $M_R$  181.88) is deposited in the flues. It is an odorless gas (Environmental Health and safety, 2005) Inhalation is the principal route of entry into the body and may result acutely in severe pneumonitis with associated mucus membrane irritation and gastrointestinal disturbances. Ambient vanadium pentoxide dust produces irritation of the eyes, nose and throat (Hauser, et al., 1995). Most absorbed vanadium is excreted in the urine within one day after a long-term moderate exposure to vanadium dust (Kiviluoto et al., 1981).

Vanadium is widely distributed in body tissues; principle organs of vanadium retention are kidneys, liver, testicles, spleen and bones. Inhalation of vanadium pentoxide can injure the lungs and bronchial airways (Environmental Health and safety, 2005) possibly involving acute chemical pneumonitis, pulmonary edema and/or acute tracheobronchitis (Nemery, 1990) Symptoms include irritation and inflammation of the mucus membranes, nasal passages and pharynx. (Environmental Health and safety, 2005). Chronic ingestion of vanadium may have significant consequences for infertility by damaging spermatogenesis. Studies in mice have demonstrated that inhalation of vanadium pentoxide results in necrosis of spermatogonium, spermatocytes and Sertoli cells (Fortoul et al., 2007). Other atmospheric pollutants will include dust generated during the maintenance works through materials handling, vehicle movements on loosely surfaced roads, transportation of uncovered construction materials (e.g. sand), and exhaust emissions from static equipment.

Workers in quarries as well as residents living around quarries are also likely to inhale dust particles into the lungs which normally results in silicosis, tuberculosis and bronchitis, which leads to pulmonary fibrosis and premature death (Decardi-Nelson *et al.*, 2015; Madhavan and Raj, 2005). However, this study revealed the concentration of vanadium and some air quality parameters in stone quarry in Abia State Nigeria.

## 2. Material and Methods

### The study area:

The study area is located in Uturu, Isuikwuato L.G.A. and Umunneochi L.G.A Abia state.

### Determination of air quality parameters

Crowcon Gasman Monitors were used for the determination of the ambient concentration of air parameters in each location from December, 2016 to July, 2017 with the months of December, January, February representing dry season and May, June July representing wet season (Asuoha and Osu, 2015). The instrument is calibrated before use and was moved from one site to the other between the hours of 8 am to 12 pm.

### Determination of vanadium pentoxide in ambient air.

Vanadium pentoxide was collected by drawing air in a stationary or personal sampler through a membrane filter made of polycarbonate, cellulose esters and/or teflon. The filter containing the collected air particulates was analyzed for vanadium using destructive method, the filter is digested in a mixture of concentrated mineral acids (hydrochloric acid, nitric acid, sulfuric acid, perchloric acid) and the vanadium concentration in the digest determined by Atomic Absorption Spectrophotometer (Kiviluoto *et al.*, 1979).

### Air Pollution Related Diseases Affecting Households

Well-structured questionnaire was administered to ninety eight residents randomly selected from the communities hosting quarries. Questions were asked on residency period in the community, perception about quarrying activities in the area and air pollution related diseases affecting households.

## 3. Results

### Result and Discussion

Concentration of SPM ranged from  $10.40 \pm 0.02$  to  $18.70 \pm 1.40 \text{ mg/m}^3$ , dry season;  $1.07 \pm 0.02$  to  $2.50 \pm 0.09 \text{ mg/m}^3$ , wet season (Table 1) for site A and  $12.01 \pm 1.10 \text{ mg/m}^3$  to  $13.60 \pm 0.30 \text{ mg/m}^3$ , wet season;  $16.40 \pm 1.30$  to  $21.30 \pm 1.60 \text{ mg/m}^3$ , dry season for site B (table 2). Highest concentration ( $18.70 \text{ mg/m}^3$ ) of SPM was observed in the dry season in site A. Concentration of SPM in the dry season is higher than in wet season because of higher production activities in dry season than that of wet season in the same location (stone quarry production plant) and also during wet season water from the atmosphere dissolves some of the particulates. These results were higher than FEPA limits. Presence of SPM may be from dust particle, gas flaring and vehicular emission or bush burning at the studied areas. Fine particles in the atmosphere provide suitable loci for various atmospheric conditions to occur. Water forms fine films which dissolves or

absorbs various contaminants presence in the atmosphere which reacts with each other. When two or more than two particles fuse higher and heavier particles are formed. Being heavier they tend to drift down and cause dry precipitation. The entire content of the atmosphere can be wash down by rain or dew, thus acid rain is caused (Osu and Asuoha, 2015).

Mean concentration of H<sub>2</sub>S ranges from 0.30 ± 0.05 to 1.70 ± 0.10 ppm, wet season; 2.06 ± 0.02 to 2.10 ± 0.01 ppm, dry season for site A and 0.60 ± 0.05 to 1.00 ± 0.10 ppm, wet season; 3.17 ± 0.02 to 1.30 ± 0.01 ppm, dry season for site B. H<sub>2</sub>S were present in all the study locations at a higher quantity. This shows that there are other contributory factors to emission of H<sub>2</sub>S such as fumes from automobile exhaust and

power generators. Results obtained was higher than results (with the range from 0.18 to 0.76) obtained by Aliyu et al., (2013) and ranged of 0.15 ± 0.05 to 0.30 ± 0.14 by Osu and Asuoha, (2015).

Mean concentration of NO<sub>2</sub> ranges from 0.50 ± 0.01 to 1.90 ± 0.03 ppm, wet season; 4.30 ± 0.01 to 7.00 ± 0.11 ppm, dry season for site A and 2.01 ± 0.02 to 3.30 ± 0.02 ppm, wet season; 4.70 ± 0.01 to 6.30 ± 0.10 ppm, dry season for site B. NO<sub>2</sub> is recorded in all study locations. Higher concentration of NO<sub>2</sub> was observed in the dry season than the wet season which is attributed to higher rate of activities in the site than the wet season and also it may be due to the use of fertilizer in agriculture.

Table 1: Concentration levels of air quality parameters in stone quarry environment A

Period	V <sub>2</sub> O <sub>5</sub> (Mg/L)	SO <sub>2</sub>	H <sub>2</sub> S	NO <sub>2</sub>	SPM (mg/m <sup>3</sup> )
December	0.18 ± 0.03	5.14 ± 0.20	2.07 ± 0.04	7.00 ± 0.11	18.70 ± 1.40
January	0.14 ± 0.01	6.10 ± 0.08	2.10 ± 0.01	4.30 ± 0.01	15.20 ± 0.60
February	0.17 ± 0.01	5.30 ± 0.06	2.06 ± 0.02	4.50 ± 0.40	10.40 ± 0.02
May	0.13 ± 0.01	0.80 ± 0.04	0.60 ± 0.01	1.90 ± 0.03	2.16 ± 0.01
June	0.13 ± 0.01	0.70 ± 0.06	0.30 ± 0.05	0.84 ± 0.10	1.07 ± 0.02
July	0.15 ± 0.01	1.20 ± 0.02	1.70 ± 0.10	0.50 ± 0.01	2.50 ± 0.09

Table 2: Concentration levels of air quality parameters in stone quarry environment B

Period	V <sub>2</sub> O <sub>5</sub> (Mg/L)	SO <sub>2</sub>	H <sub>2</sub> S	NO <sub>2</sub>	SPM (mg/m <sup>3</sup> )
December	0.46 ± 0.01	6.20 ± 0.10	3.17 ± 0.02	6.30 ± 0.10	21.30 ± 1.60
January	0.36 ± 0.02	6.80 ± 0.04	3.10 ± 0.07	5.12 ± 0.06	19.20 ± 0.60
February	0.18 ± 0.02	6.40 ± 0.40	1.30 ± 0.01	4.70 ± 0.01	16.40 ± 1.30
May	0.11 ± 0.02	5.10 ± 0.15	1.00 ± 0.01	3.30 ± 0.02	13.60 ± 0.30
June	0.10 ± 0.01	1.30 ± 0.02	0.60 ± 0.05	2.01 ± 0.02	13.11 ± 0.70
July	0.11 ± 0.03	1.60 ± 0.03	0.90 ± 0.01	2.10 ± 0.01	12.01 ± 1.10

Table 3: Concentration levels of heavy metals in dust of stone quarry A

Period	Cu	Cr	Zn	Hg	Pb	Ni
December	42.50	38.70	136.30	0.65	16.80	16.40
January	31.80	32.30	98.80	0.80	13.40	14.50
February	29.90	23.50	91.00	0.50	11.06	8.10
May	18.90	34.70	74.70	0.10	15.10	5.10
June	14.40	25.60	45.60	ND	11.80	2.00
July	14.00	25.80	48.80	ND	10.60	1.60

Table 4: Concentration levels of heavy metals in dust of stone quarry B

Period	Cu	Cr	Zn	Hg	Pb	Ni
December	25.10	15.20	186.10	0.20	13.90	5.40
January	19.30	11.80	128.70	0.05	11.20	4.10
February	14.60	9.90	101.00	0.01	9.00	2.10
May	13.90	4.70	91.40	ND	11.10	1.60
June	13.00	5.90	74.30	ND	8.80	1.08
July	11.30	7.70	44.30	ND	6.60	1.00

Mean concentration of SO<sub>2</sub> in study locations shows the same trend of being higher during dry season and lower concentration during wet season. This is due to dilution effect of rain on the atmosphere. Its values ranges from 0.70 ± 0.06 to 1.20 ± 0.02 ppm, wet season; 5.14 ± 0.20 to 6.10 ± 0.08 ppm, dry season for site A and 1.30 ± 0.02 to 5.10 ± 0.15 ppm, wet season 6.20 ± 0.10 to 6.80 ± 0.04 ppm, dry season for site B. Higher values during dry season may be attributed to vehicular emissions, refuse dump and other combustion activities within the area. SO<sub>2</sub> is known as an acid precursor and was identified in all samples studied. The concentration of SO<sub>2</sub> is lower than ranges of 3.21 to 5.18 ppm, 7.4 to 15.5 ppm, and 16 to 64 ppm reported by Ettouney et al. (2010), and Kalabokas et al. (1999), respectively but was higher than range of 0.03 to 0.09 ppm reported in Kano metropolis, Nigeria (Okunola et al., 2012) and range of 0.022 to 0.087 ppm reported by (Aliyu et al., 2013), Osu and Asuoha (2015) with range of 0.95 ± 0.06 to 2.00 ± 0.11 ppm. Comparing data with the FEPA

standard level of 0.06 ppm, the results obtained from this study was found to be higher.

Vanadium is a redox-sensitive metal that is released to soils by weathering and anthropogenic emissions. Vanadium is released naturally to the atmosphere by the formation of continental dust, marine aerosols, and volcanic emissions. Vanadium is a constituent of nearly all coal and petroleum crude oils. The concentration of vanadium in the environment ranged from  $0.13 \pm 0.01$  to  $0.18 \pm 0.03$  and  $0.10 \pm 0.01$  to  $0.46 \pm 0.01$  for sample A and B respectively. Combustion of heavy fuels, especially in oil-fired power plants, refineries, and industrial boilers, and coal are the major source of anthropogenic emissions of vanadium into the atmosphere (Mamane and Pirrone 1998; Sepe et al. 2003). During the combustion of residual oils organovanadium compounds found in fuel oils are oxidized and transformed into various compounds (e.g., vanadium pentoxide, vanadium tetroxide, vanadium trioxide, and vanadium dioxide). These compounds are emitted as fly ash into the atmosphere (Mamane and Pirrone 1998).

Vanadium is the 22<sup>nd</sup> most abundant element in the earth's crust (Baroch 2006). There are about 65 different vanadium-containing minerals; carnotite, roscoelite, vanadinite, and patronite are important sources of this metal along with bravoite and davidite (Baroch 2006; Lide 2008). It is also found in phosphate rock and certain ores and is present in some crude oils as organic complexes (Lide 2008).

Distribution of Pb, Hg, Cu, Zn, Cr and Ni in dust from stone quarry environment were shown in tables 3 and 4. The concentration of the heavy metal ranged from 14.00 to 42.50 mg/kg, Cu; 23.50 to 38.70 mg/kg, Cr; 45.60 to 136.30 mg/kg, Zn; 0.10 to 0.65 mg/kg, Hg; 10.60 to 16.80 mg/kg, Pb and 1.60 to 16.40 mg/kg, Ni for site A. Concentration of the heavy metal in site B ranged from 11.30 to 25.10 mg/kg, Cu; 4.70 to 15.20 mg/kg, Cr; 44.30 to 186.10 mg/kg, Zn; 0.01 to 0.20 mg/kg, Hg; 6.60 to 13.90 mg/kg, Pb and 1.00 to 5.40 mg/kg, Ni. Concentration of lead is high in the studied samples. The dust samples showed significant higher values of the heavy metal contents. This may be due to the quarry activities and accumulation of the automobile wastes in the sampled soils.

The health problems suffered by the residents are those that associate with inhalation of dust in the air. According to reports, the detonation of explosives in quarrying operation causes ground vibration, which produces effects such as stress, anxiety, and increased pulse rate, loss of sleep, fatigue and excessive contraction of pupil among residents living near quarry site (Salvato, 1992., Alloway and Ayres, 1995). This study found that about 99.99 % of air pollution diseases affect the households. The workers

had no safety measures. Data from this study showed that chronic exposure to dust and polluted air from crushing of stones increased susceptibility to respiratory problems and increased length of service is an additional risk factors. Percentage values obtained were 24.49 %, catarrh; 28.57 %, coughing; 26.53 %, headach; 9.18 %, chest pains and 11.22 %, eye infections. Nwibo *et al.*, (2012) found out that the workers had various respiratory problems including chest pain, cough, wheezing and shortness of breath. Ugbogu, *et al.*, (2009) studied the occurrence of respiratory and skin problems among manual stone workers and found out that up to 85% of the workers had respiratory symptoms while 77% had skin infection. Ilyas *et al.*, (2010) established that dust related problems were exacerbated by cases of owners of the crushing stone unit not providing appropriate measures to protect the workers. the health problems experienced by quarry workers are identical, and are similar to those identified by Olusegun et al.,2009.

### Conclusion

Stone quarrying has been shown as a human activity that generates particulates pollution in the environment. Varying concentrations of air contaminants were detected at all sampling points. The high level of particulates generated at the drilling and crushing areas depicts them as hazard zones. Higher concentration of air parameters were observed in the dry season than wet season. Moreover, exposures of quarry workers to air contaminants dispose them to several health problems (respiratory ailments) which is common among residents living near quarry sites. Considering the various prevailing health problems associated with inhalation of air contaminants, environmental impact assessment should be conducted for all quarries.

### Corresponding Author:

Osu Charles I

Department of Pure and Industrial Chemistry,  
University of Port Harcourt, P.M.B 5323 Choba, Port  
Harcourt, Rivers State, Nigeria.

Telephone: +234 803 7783246

E-mail: [charsike@yahoo.com](mailto:charsike@yahoo.com)

### References

1. Bauer, G., Güther, V., Hess, H., Otto, A., Roidl, O., Roller, H. & Sattelberger, S. (2003) Vanadium and vanadium compounds. In: Bohnet *et al.*, eds, *Ullmann's Encyclopedia of Industrial Chemistry*, 6th rev. Ed., Vol. 38, Weinheim, Wiley-VCH Verlag GmbH & Co., pp. 1–21.
2. Lide, D.R., ed. (2003) *CRC Handbook of Chemistry and Physics*, 84th Ed., Boca Raton, FL, CRC Press LLC, p. 4-93.

3. Kucera, J. & Sabbioni, E. (1998) Baseline vanadium levels in human blood, serum, and urine. In: Nriagu, J.O., ed., *Vanadium in the Environment. Part 2: Health Effects*, New York, John Wiley & Sons, pp. 75–89.
4. Kiviluoto, M., Pyy, L. & Pakarinen, A. (1979) Serum and urinary vanadium of vanadium-exposed workers. *Scand. J. Work Environ. Health*, 5, 362–367.
5. National Toxicology Program (2002) *Toxicology and Carcinogenesis Studies of Vanadium Pentoxide (CAS No. 1314-62-1) in F3344/N rats and B6C3F1 mice (Inhalation Studies)* (Technical Report series No. 507; NIH Publication No. 03-4441) Research Triangle Park, NC.
6. Ress, N.B., Chou, B.J., Renne, R.A., Dill, J.A., Miller, R.A., Roycroft, J.H., Hailey, J.R., Haseman, J.K. & Bucher, J.R. (2003) Carcinogenicity of inhaled vanadium pentoxide in F344/N rats and B6C3F1 mice. *Toxicol. Sci.*, 74, 287–296.
7. O'Neil, M.J., ed. (2001) *The Merck Index*, 13th Ed., Whitehouse Station, NJ, Merck & Co., Inc., p. 1767.
8. Fortoul TI, Bizarro-Neves P, Acevedo-Nava S, Piñón-Zárate G, Rodríguez-Lara V, Colin-Barenque L, et al. Ultrastructural findings in murine seminiferous tubules as a consequence of subchronic vanadium pentoxide inhalation. *Reprod Toxicol.* 2007;23:588–92. [PubMed].
9. Nemery B. Metal toxicity and the respiratory tract. *Eur Respir J.* 1990;3:202–19. [PubMed].
10. Environmental Health and Safety; (2005). Vanadium pentoxide. Phillipsburg, New Jersey: MSDS; p. 5.
11. Hauser R, Elreedy S, Hoppin JA, Christiani DC. Upper airway response in workers exposed to fuel oil ash: nasal lavage analysis. *Occup Environ Med.* 1995;52:353–8. [PubMed].
12. Kiviluoto M, Pyy L, Pakarinen A. Serum and urinary vanadium of workers processing vanadium. *Int Arch Occup Envir Health.* 1981;48:251–6. [PubMed].
13. Wu, C., Lin, M., Feng, C., Yang, K., Lo, Y. and Lo, J. (2003). Measurement of Toxic Volatile Organic Compounds in Indoor Air of Semiconductor Foundries Using Multi-sorbent Adsorption/ Thermal Desorption coupled with Gas Chromatography-Mass Spectrometry. *Journal of Chromatography Vol. 996:* 225-231.
14. Aliyu, Y. A., Musa, I. J, Youngu, T. T. (2013). Appraisal of Sulphur Contaminants from Transportation in Urban Zaria, Nigeria. *Academic Journal of Interdisciplinary Studies MCSER Publishing, Rome-Italy Vol 2 No 10: 155 -163.*
15. Osu Charles I and Asuoha, Adaku N. (2016) Comparative Study Of The Prevailing Illness In Ibeno And Its Environs In Akwa Ibom State Linked To Overload Of Air Quality Parameters. *Researcher* 2016;8(6): 42 – 49.
16. Kalabokas, P. O., Viras, L. G. and Repapis, C. (1999), “Analysis of the 11 – years record (1987 - 1997) of Air Pollution Measurements in Athens, Greece, Part I: Primary Air Pollutants”, *Global Nest Journal*, Vol. 1 No. 3, pp. 157 – 167.
17. Ettouney, R. S., Zaki, J. G., El-Rifiai, M. A. and Ettouney, H. M. (2010), An Assessment of the Air Pollution Data from Two Monitoring Stations in Kuwait, *Toxicology and Environmental Chemistry*, Vol. 92 No. 4, pp. 655 – 668.
18. Okunola, O. J., Uzairu, A., Gimba C. E. and Ndukwe, G. I. (2012). Assessment of Gaseous Pollutant along High Traffic Roads in Kano, Nigeria. *International Journal of Environment and Sustainability*. Volume 1(1): 1-15.
19. Madhavan., P. and Raj. S, (2005). Budhpura ‘Ground Zero’ Sandstone quarrying in India. *Mine Labour Protection Campaign.*
20. Isaac Decardi-Nelson, Betty Solomon-Ayeh, Daniel Asenso-Gyambibi, Kwadwo Twumasi-Ampofo\*,. and Alex Ahenkuro. (2015). An Assessment of the Institutional Capacity for the Management of Quarry Industries in Ghana: The Case of Buoho Stone Quarries. *Civil and Environmental Researc.* Vol.7, No.5., 84 – 97.
21. Oguntoke Olusegun, Aboaba Adeniyi and Gbadebo T. Adeola (2009). Impact of Granite Quarrying on the Health of Workers and Nearby Residents in Abeokuta Ogun State, Nigeria. *Ethiopian Journal of Environmental Studies and Management Vol.2 No.1.* 1 – 11.
22. Baroch, F., 2006. Vanadium and vanadium alloys. *Kirk-Othmer Encyclopedia of Chemical Technology*pp. 1–18.
23. Nriagu, J., Pirrone, N., (1998). Emission of vanadium into the atmosphere. In: Nriagu, J. (Ed.), *Vanadium in the Environment. Part I: Chemistry and Biochemistry*. Wiley, New York, pp. 25–36.
24. Alloway, B. J. and Ayres, D. C. (1995). *Chemical Principles of Environmental Pollution*. Blackie Academic and Professional, London. Pg. 250-290.
25. Ugboogu, O.C., Ohakwe, J. & Foltescu, V. (2009). Occurrence of respiratory and skin problems among manual stone-quarrying workers. *African Journal of Respiratory Medicine*, March, 2009 pp 23-26.

26. Salvato, J. A. (1992). *Environmental Engineering and Sanitation*. 4th Edition – John Wiley and Sons Incorporated. Pg. 767- 806.
27. Ilyas, M. & Rasheed, F. (2010). Health and environment related issues in stone crushing in Pakistan. Paper submitted on 10th Round Regional Research Competition of South Asia Network of economic Research Institute (Sanei).
28. Nwibo, A.N., Ugwuja, E.I., Nwambeke, N.O., Emelumadi, O.F. & Ogonnaya, L.U. (2012). Pulmonary problems among quarry workers of crushing stone industrial site at Umuoghara, Ebonyi state, Nigeria. *The International Journal of Occupational and Environmental Medicine* 2012; 3: 178-185.
29. Mamane, Y. and Pirrone, N. (1998). *Vanadium in the atmosphere*. In: Part 1: Chemistry and Biochemistry, New York, John Wiley and Sons. pp. 37–71.
30. Sepe, A., Ciaralli, L., Ciprotti, M., Giordano, R., Funari, E. and Costantini, S., (2003). Determination of cadmium, chromium, lead and vanadium in six fish species from the Adriatic Sea. *Food Addit. Contam.* 20: 543–552.

10/23/2017