Wound healing Benefits of Curcumin for Perineal Repair after Episiotomy: Results of an Iranian Randomized Controlled Trial

Safar Ali Esmaeili Vardanjani¹, Fahimeh Sehati Shafai², Parvin Mohebi³, Marjan Deyhimi⁴, Abbas Delazar⁵ Morteza Ghojazadeh⁶, Padideh Malekpour⁷

 ^{1.} Msc in Nursing Education, Shahrekord University of Medical Science, Shahrekord, Iran.
^{2.} MSc in midwifery, Faculty Member of Nursing and Midwifery School, Tabriz University of Medical Science, Tabriz, Iran.

^{3.} MSc in Midwifery, Zanjan University of Medical Science, Zanjan, Iran.

⁴ MSc in midwifery, International Branch of Shahid Beheshti University of medical science. Tehran, Iran.

⁵ PhD in pharmacogenosy, associated professor of pharmacy faculty of Tabriz medical science university.

⁶ PhD In Physiology, assistant professor of medical faculty, Tabriz medical science university.

⁷ MSc in midwifery, Drug Applied Research Center, Tabriz University of Medical science, Tabriz, Iran.

padideh_66@yahoo.com

Abstract: Pain and discomfort related to episiotomy have been reported to interfere with women's daily activities postpartum, such as sitting, walking and lifting the baby. To compare the effects of curcumin and Povidone-iodine solutions for episiotomy healing in primiparous women. 120 healthy primiparous women with a vaginal delivery at term were evaluated in this double-blind randomized clinical trial. Randomization was done using a table of random list numbers. Perineal healing was evaluated by research midwives blinded to random allocation at 24–48 hours and 10 days postpartum. Pain was assessed via a visual analogue scale and wound healing via the REEDA scale. Analysis was done on the intention-to-treat principle. The main outcome measure was the changes in wound healing between the two groups as measured by the REEDA Scale. Secondary outcome measures were perineal pain and wound healing 24–48 hours and 10 days after delivery. There was a greater decrease in the total scores of the REEDA in the curcumin group than in the Povidone-iodine (P < 0.001), however; there weren't significant differences between the groups on the VAS scores. The application of curcumin may assist in the episiotomy healing process and could be suitable replacement for Povidone-iodine.

[Esmaeili Vardanjani SA, Sehati Shafai F, Mohebi P, Deyhimi M, Delazar A, Ghojazadeh M, Malekpour M. Wound healing Benefits of Curcumin for Perineal Repair after Episiotomy: Results of an Iranian Randomized Controlled Trial. *Life Sci J* 2012;9(4):5536-5541] (ISSN:1097-8135). http://www.lifesciencesite.com. 821

Keywords: Episiotomy, Curcumin, Povidone-Iodine, Perineal Pain, Wound Healing.

1. Introduction

Perineal trauma is a frequent complication to vaginal delivery, and more than 90% of primiparous women in developing countries sustain episiotomy (Morhe et al., 2004).

Episiotomy's pain and discomfort have been related to difficulties in women's daily activities in postpartum, such as sitting, walking and lifting the baby. (Albers et al., 1999). Pain related to episiotomy is identified to have a negative impact on sexual activities in the first year after childbirth (Glazener, 1997). Midwives and obstetricians increasingly face women who wish to have a caesarean section due to fear of genital tract injuries or following previous childbirth trauma (McCourt et al 2007; Wagner, 2000).

Reducing perineal trauma and associated morbidity has a high importance for childbearing women and health professionals (Homer and Dahlen ,2007). Some existing treatments for episiotomy healing aren't effective and some may prolong the healing processes (Steen et al , 2000).

Povidone-iodine is an antiseptic solution that is usually used in Iran for episiotomy healing. (Vakilian et al., 2011). Tork and Valaei (2002) demonstrated that there was no significant difference between the Povidone-iodine and water in episiotomy wound healing. Cooper et al (1991) even showed that Povidone-iodine suppresses function of fibroblasts and lymphocytes.

Alternative and complementary medicine is used widely in dermatologic surgery such as episiotomy healing. (Reddy et al., 2011) Turmeric is considered as the foundation of an herbal programmed for health. For thousands of years it has been believed to be a key balancing and detoxifying herb. The main biologically active part of Turmeric is curcumin. Curcuminoids are natural phenols that are responsible for the yellow color of turmeric. (Redfern R. 2004). Previous research suggested that curcumin has anti-inflammatory and pro-wound-healing effects and it exerted its anti-inflammatory effects (Sidhu GS et all 1998).by increasing the mRNA transcripts of growth factor-beta 1 and fibronectin .It is notice worthy that only few number of randomiezed controlled trials have been conducted in humans to demonstrate its effective properties (White and Judkins, 2011; Reddy et al., 2011).

Therefore, this study was performed to compare the effects of curcumin and Povidone-iodine solutions for episiotomy healing in primiparous women.

2. Material and Methods

The prospective, parallel randomized clinical trial was conducted with recruitment through 2009 -2010 in the Alzahra Teaching Hospital in Tabriz, Iran. The study protocol was approved by the Ethics Committee and research center of the Tabriz University of Medical Sciences. (14.89.9 No). The trial was registered at the Iranian Registry of Clinical Trials at <u>www.irct.ir</u> (IRCT138810193027N1).

Eligibility and recruitment

Eligible subjects were primiparous women, without any acute and chronic disease or allergy, and had a healthy pregnancy and delivery after 37 weeks of gestation. The women also had to be able to read and speak in native language. Exclusion criteria were perineal injuries involving the anal sphincter and/or anal mucosa, postpartum hemorrhage, previous perineal surgery, obvious violence and sexual intercourse in the early postpartum period.

Written informed consent was obtained from all women who decided to participate in the study. Participants not meeting the eligibility criteria but randomized included in the analysis according to the intention-to-treat principle.

Randomization

In this study, 120 patients participated in research project .We prepared 60 Curcumin solution as well as 60 Povidone -iodine solution tubes. It is notice worthy that all 120 tubes were alike and each tube had its specific code. Therefore, researcher and patients didn't inform of the content of tubes. Participants consumed Curcumin solution categorized as an intervention group, participants who consumed the Povidone-iodine solution were as a control group (60 subjects in each arm). Therefore, each patient had a specific tube and she should consume it three times a day. It is notice worthy those women must wash their hands before using solutions in order to reach acceptable results, and every necessary information related to consuming solutions were explained to each participant individually.

Scales and Outcome of study

REEDA scale is as a descriptive scale, it can measure five components associated with healing process ,they are including :Redness ,Edema ,Ecchymosed, Discharge , Approximation ,and it is graded 0-3 .Therefore ,each category is evaluated and a number assigned for a total REEDA score ranging from 0-15.In this study ,REEDA scale was applied to represent the changes in wound healing among two solutions. Consequently, REEDA scores for curcumin and Povidone-iodine groups were 2.5 ± 0.8 and 3.7 ± 1.6 respectively.

The Visual Analogue Scale (VAS) is a subjective measure of pain. It consists of a 10cm line with two end-points representing 'no pain' and 'Unbearable pain'. Patients are asked to rate their pain by placing a mark on the line corresponding to their current level of pain According to this method, score 0 (No pain), 1-3 (Mild pain), 4-7(Moderate pain), and 8-10 (Unbearable pain). Consequently, Pain was evaluated by VAS method in our research.



Both Curcumin and Povidone-iodine solutions were packed in similar cans containing a code which was known only to the study pharmacist. The study was double-blinded and the contents were known neither to the participants, nor to the research midwives

Statistical analysis

All statistical analyses were done on an intention-to-treat basis. Data were entered twice into the software SPSS 16 software to correct for typing errors. Continuous data with normal distribution were analyzed by independent two-sample *t*-tests and without normal distribution by Mann–Whitney U test. For categorical variables the Chi-Square and Fisher's exact tests were used. All statistical tests were two-tailed, and the significance level was set at 0.05 ($\alpha = 0.05$). Values are described as the mean (standard deviation) or the median (interquartile range) if not normally distributed.

3. Results

The CONSORT diagram shows the participant flowchart through the trial (Figure 1). Three women withdrew their consent to participate. Thus, the number of intervention and Control group were 59 and 58 respectively.



Figure 1 Consort diagram of vaginal deliveries in women during study period

Table 1 shows baseline characteristics of the curcumin and Povidone-iodine groups. The two Groups were comparable and did not differ with respect to any of the baseline demographics and delivery details variables, indicating successful randomization.

Table 2 represents the comparison of episiotomy healing evaluation in the two groups on basis of REEDA scale. As shown in table 2, at 24 hours 8 participants in the curcumin group and 0 of the Povidone-iodine group had ecchymosis. Within 0.25 cm, which was significantly different between the groups (P = 0.006). Within 10 days more

participants in the curcumin group than the Povidoneiodine group had redness (P = 0.033) and edema (P = 0.027). The other REEDA parameters were not significantly different between the groups.

According to Table .3, The REEDA scores were significantly differed between the both groups at 24 hours (P = 0.032) and 10 days (P < 0.001). The VAS score differences between the two groups were statistically not significant at 24 hours (P = 0. 2) and significant in 10 days (P < 0.001); however, difference scores between 10^{th} and first day were not statically significant.

٦

Table 1 Characteristics of the two groups at baseline							
Variables	Curcumin group $(N = 59)$	Povidone - Iodine group ($N = 58$)	P value				
Age (years)	24.2 ± 5.6	22.7 ± 4.0	0.08				
Body mass index (kg/m ²)	25.8 ± 3.6	26.1 ± 3.0	0.58				
Gestational age (weeks)	38.1 ± 5.1	38.6 ± 0.8	0.46				
Education (diploma)	28 (47.5)	21 (36.2)	0.35				
Job (housewife)	56 (94.9)	49 (84.5)	0.61				
Status of operator (supervising midwife)	20 (33.9)	11 (19)	0.61				
Note: Data are presented as Mean ± SD (standard deviation) and Frequency (Percentage).							

Table 2. Comparison	of episiotomy healing evaluation	on in the	e two groups			
Variables			Curcumin group (N =	Povidone-iodine	P value	
			59)	group ($N = 58$)	1 value	
REEDA Parameters a	t 24 hours					
Redness	T					
	None		0 (0)	2 (3.4)	0.244	
	0.25 cm or more		59 (100)	56 (96.6)		
Edema	1					
	None		6 (10.2)	2 (3.4)	0.272	
	≤ 1 or more		53 (89.8)	56 (96.6)		
Ecchymosis						
	None		35 (81.4)	37 (100)	0.006	
	0.25 cm or more		8 (18.6)	0 (0)		
Discharge						
	None to serum		53 (89.8)	57 (98.3)	0.114	
	Serosanguinous or bloody		6 (10.2)	1 (1.7)		
Approximation						
	Closed		53 (89.8)	57 (98.3)	0.114	
	Separation <= 3 mm or more	•	6 (10.2)	1 (1.7)		
REEDA Parameters a	t 10 days					
Redness						
	None		38 (64.4)	26 (44.8)	0.033	
	0.25 cm or more		21 (35.6)	32 (55.2)		
Edema						
	None $\leq 1 \text{ cm}$		50 (84.7)	39 (67.2)	0.027	
			9 (15.3)	19 (32.8)	0.027	
Ecchymosis	•		· · ·			
	None		58 (98.3)	57 (98.3)	1	
	0.25 cm or more		1 (1.7)	1 (1.7)		
Discharge	•					
	None to serum		49 (83.1)	45 (77.6)	0.457	
	Serosanguinous or bloody		10 (16.9)	13 (22.4)	0.457	
Approximation						
	Closed		55 (93.2)	49 (84.5)	0.122	
	Separation <= 3 mm or more		4 (6.8)	9 (15.5)	0.133	
	Note: Data are prese	ented as	Frequency (Percentag	je).	1	
	1					
Т	able 3 Comparison of the two (Groups a	at 24 hours and 10 da	ys after deliver		
Variables	•	Curcu	min group (N=59)	Povidone-iodine group (N=58)	P value	
VAS score		1		0 ··· r (· · · · · /		
	At 24 hours	5 (4,5))	5 (4,5)	0.027	

	At 10 days	0 (0,0)	1(0,1)	< 0.001		
	Difference scores between 10 th and first day	-4 (-5,-4)	-4 (-5,-4)	0.963		
REEDA score						
	At 24 hours	5 (4,5)	5 (4,5)	0.032		
	At 10 days	0 (0,1)	1 (1,2)	< 0.001		
	Difference scores between 10 th and first day	-4 (-5,-4)	-3 (-3,-2)	< 0.001		
Note: Data are presented as Median (Interquartile Range).						

A per-protocol analysis was performed to investigate whether the noncompliance had contributed to validating a false difference between the two groups. The per-protocol analysis showed statically significant differences in the first outcome measure over again (Ps < 0.05, data not shown). There were no effects for the curcuminand Povidone-iodine groups.

4. Discussions

The main aim of this study was evaluating the role of Curcumin in wound healing in compare to Povidone-iodine .Our results showed that the curcumin solution for perineal episiotomies was more effective way to improvement of wound healing processes than the Povidone-iodine. However curcumin wasn't effective in pain relief.

Wound healing is a dynamic, interactive process concerning soluble mediators, blood cells, extracellular matrix, and parenchymal cells. Wound healing has at least three phases -inflammation, proliferation, and maturation - which overlap in time. (Singer and Clark, 1999).

Treatment with curcumin could enhance the synthesis of collagen, DNA, fibroblast, vascular densities and other important factors in wound healing. (Ghasemi Dehkordi et al (2003) demonstrated that curcumin pretreatment had a helpful effect on the irradiated wound and could be a substantial therapeutic strategy for improving radiation induced delay in wound repair in cases of radiation-induced skin injuries. Also Sidhu et al (1999) showed that curcumin accelerate wound healing in diabetic rats by increased formation of granulation tissue, faster reepithelialization and increased collagenization.

The results of a small clinical trial with 19 HIV patients found that a liquid soap of curcumin could decrease itching symptoms and infectious wound and abscess. (Hong et al., 2004). This study, as of first clinical trials conducted in humans showed that curcumin solution had wound healing effects.

Curcumin solution wasn't effective in episiotomy pain relief in this study. Vakillian et al (2011) also showed that lavender essential oil could not cause to decrease of pain. These studies may highlight the need for future trials that can employ interventions of greater strength of the clinician– patient relationship because pain relief is never about the clinician's intervention alone.

Limitations of the present study might be related to the absent of control on the nutrition and the stresses that may contribute to wound healing and pain relief. The authors believe that future trials could consider these issues.

It seems that curcumin can be used as an appropriate treatment for postpartum episiotomy wound care according to the familiarity of the mothers with complementary medicine. Consequently, the authors suggest the assessment of its application in other areas such as cesarean section and umbilical wound healing and also considering the use of curcumin solution instead of Povidone-iodine for episiotomy wound care.

Acknowledgements:

This study was performed with financial support from Drug Applied Research Center, Tabriz University of Medical science, Tabriz, Iran. Also, authors express their appreciation for all participants of this study that they made this study possible.

Corresponding Author:

Padideh Malekpour

MSc in midwifery, Drug Applied Research Center Tabriz University of Medical science, Tabriz medical science university, nurse-midwifery faculty, Tabriz iran.

E-mail: *padideh 66@yahoo.com*

References

- 1. Morhe ES, Sengretsi S, Danso KA. Episiotomy in Ghana. Int J Gynaecol Obstet 2004;86(1):46-7.
- Albers L, Garcia J, Renfrew M, McCandlish R, Elbourne D. Distribution of genital tract trauma in childbirth and related postnatal pain. Birth 1999;26(1):11-7.
- 3. Glazener CM. Sexual function after childbirth: women's experiences, persistent morbidity and

lack of professional recognition. Br J Obstet Gynaecol 1997;104(3):330-5.

- 4. McCourt C, Weaver J, Statham H, Beake S, Gamble J, Creedy DK. Elective cesarean section and decision making: a critical review of the literature. Birth 2007;34(1):65-79.
- 5. Wagner M. Choosing caesarean section. Lancet 2000;356(9242):1677-80.
- Homer C, Dahlen H. Obstetric-induced incontinence: a black hole of preventable morbidity? An 'alternative' opinion. Aust N Z J Obstet Gynaecol 2007;47(2):86-90.
- Steen M, Cooper K, Marchant P, Griffiths-Jones M, Walker J. A randomised controlled trial to compare the effectiveness of ice-packs and Epifoam with cooling maternity gel pads at alleviating postnatal perineal trauma. Midwifery 2000;16(1):48-55.
- Vakilian K, Atarha M, Bekhradi R, Chaman R. Healing advantages of lavender essential oil during episiotomy recovery: a clinical trial. Complement Ther Clin Pract 2011;17(1):50-3.
- 9. Tork ZS, Valaei N. Comparison of the effect of Betadine and water in episiotomy wound healing. FEYZ 2002.
- 10. Cooper ML, Laxer JA, Hansbrough JF. The cytotoxic effects of commonly used topical antimicrobial agents on human fibroblasts and keratinocytes. J Trauma 1991;31(6):775-82.
- 11. Reddy KK, Grossman L, Rogers GS. Common complementary and alternative therapies with potential use in dermatologic surgery: Risks and benefits. J Am Acad Dermatol 2011.
- 12. Redfern R. Pain Relief, Inflammation Relief and Clear Arteries! 2004.
- 13. White B, Judkins DZ. Clinical Inquiry. Does turmeric relieve inflammatory conditions? J Fam Pract 2011;60(3):155-6.
- 14. Ghasemi Dehkordi N, Sajjadi SE, Ghannadi AR. Iranian herbal pharmacopoeia (IHP). Hakim Research Journal 2003;3(6):69-3.
- 15. Davidson N. REEDA: evaluating postpartum healing. J Nurse Midwifery 1974;19(2):6-8.

- 16. Moher D, Schulz KF, Altman DG. The CONSORT statement: revised recommendations for improving the quality of reports of parallel group randomized trials. BMC Med Res Methodol 2001;1:2.
- 17. Singer AJ, Clark RA. Cutaneous wound healing. N Engl J Med 1999;341(10):738-46.
- Jagetia GC, Rajanikant GK. Role of curcumin, a naturally occurring phenolic compound of turmeric in accelerating the repair of excision wound, in mice whole-body exposed to various doses of gamma-radiation. J Surg Res 2004;120(1):127-38.
- Sidhu GS, Mani H, Gaddipati JP, Singh AK, Seth P, Banaudha KK, Patnaik GK, Maheshwari RK. Curcumin enhances wound healing in streptozotocin induced diabetic rats and genetically diabetic mice. Wound Repair Regen 1999;7(5):362-74.
- Hong J, Bose M, Ju J, Ryu JH, Chen X, Sang S, Lee MJ, Yang CS. Modulation of arachidonic acid metabolism by curcumin and related beta-diketone derivatives: effects on cytosolic phospholipase A(2), cyclooxygenases and 5-lipoxygenase. Carcinogenesis 2004;25(9):1671-9.
- 21. Kettle C, Hills RK, Ismail KM. Continuous versus interrupted sutures for repair of episiotomy or second degree tears. Cochrane Database Syst Rev 2007;4.
- 22. Kettle C, Tohill S. Perineal care. Clin Evid (Online) 2008;2008.
- 23. Pizzo PA, Clark NM. Alleviating suffering 101--pain relief in the United States. N Engl J Med 2012;366(3):197-9.
- 24. Sidhu GS, Singh AK, Thaloor D, Banaudha KK, Patnaik GK, Srimal RC, Maheshwari RK. Enhancement of wound healing by curcumin in animals. Wound Repair Regen. 1998 Mar-Apr;6(2):167-77.

12/21/2012