Socioeconomic status and quality of life in elderly people in rural area of Sari-Iran

Samad Rouhani¹, Parsa Zoleikani²

¹ Assistant Professor, Department of Public Health; Psychiatry&Behavioral Sciences Research Center, Mazandaran University of Medical Sciences, Sari, Iran. Hospital Management Research Center, Tehran University of Medical Sciences, Tehran, Iran.

Corresponding author: parsa.z123@yahoo.com

Abstract: There are many reports particularly from developed countries that show a relationship between socioeconomic status and health of individuals. Among a wide range of socioeconomic variables it seems that income, education, and occupation are more important factors that might change health of people. Many of studies carried out to assess the relationship between socioeconomic status and health status focused on narrow health indicators that measure a single feature of health status such as morbidity and mortality. In this study we employed a specific instrument of quality of life measurement (LEIPAD) to measure the quality of life of elderly people. It was a cross-sectional study carried out in rural area of Sari the capital city of Mezandaran province. A sample of 130 elderly people over 60 years from both sex were interviewed on their doorsteps randomly. Inferential statistics and ANOVA were used to analyze the data using SPSS software package. The quality of life of respondents in overall was better on core scales than moderator scales. Among domains of core scales, physical functioning scale and social functioning scale had better status. Among socioeconomic variables age, marital status, income and literacy had significantly affected the quality of life of elderly people both in terms of core and moderator scales. It is recommended that appropriate intervention needs to assist vulnerable groups particularly elderly people in rural area to improve the quality of their life.

[Rouhani S, Zoleikani P. Socioeconomic status and quality of life in elderly people in rural area of Sari-Iran. *Life Sci J* 2013;10(2s):74-78] (ISSN:1097-8135). https://www.lifesciencesite.com. 12

Keywords: Socioeconomic; elderly people; quality of life; rural area

1. Introduction

There are many reports particularly from developed countries that show a relationship between socioeconomic status and health of individuals (Banks et al., 2006; Bassuk et al., 2002; Marmot, 2004) indicating a better health for higher socioeconomic status compared with lower socioeconomic group. Among a wide range of socioeconomic variables it seems that income, education, and occupation are more important factors that might change health of people (Lynch et al., 2000; Mustard et al., 1997). This happen because different health behavior, range of obesity, and inequality in medical treatment, poor perceived health status of different socioeconomic groups can vary their mortality rate (Bassuk et al., 2002). In developing countries. in particular, studies that show socioeconomic variables and health status are relatively limited. This limitation is more evident when it comes to specific vulnerable groups such as elderly people (Qin, 2007).

A series of theories have been proposed to explain the influences of personal and socio-environmental factors on subjective attitudes toward aging that most of them focus on the individual's adaptability to aging. Political economy perspective highlights the socio-structural influences on aging. It

emphasizes the relationship between socioeconomic determinants and older people's quality of life. It argues that social resources are distributed unequally on the basis of gender, class, and race due to political and economic forces. For instance, in the social context of developing countries in particular, female and rural older persons are less likely to have pension and public medical care and consequently, they are economically disadvantaged and are more likely to have bad quality of life. Political economy perspective also allows policy-makers to offer policy interventions to improve quality of life (QOL) of older adults (Qin, 2007).

Population aging is a global issue that draws attention from academic, political, and economic fields. It not only is characterized as a social issue in developed countries where it has been most prevalent, but also recognized as a social problem faced by more and more developing nations. Therefore, research in this area could be important for developing countries such as Iran as the findings of such investigations could provide invaluable evidence for policymakers and stakeholders to boost their infancy social security programs in dealing with vulnerable groups.

Many of studies carried out to assess the relationship between socioeconomic status and health status focused on narrow health indicators that measure

² Student Research Committee; Health Staff of Sari District Health Center, Mazandaran University of Medical Sciences, Sari, Iran

a single feature of health status such as morbidity and mortality (Huguet et al., 2008). In recent years experts recommend the use of more comprehensive indicator of health status measurement that are known as Health Related Quality Of Life (HRQQL) measures that can show health status and functional level of individuals. The World Health Organization's Quality of Life (WHOQOL) Group has defined the concept of QOL as "an individual's perception of their position in life, in the context of the culture and value systems in which they live and in relation to their goals, expectations, standards and concerns" (WHOQOL-Group, 1993). Accordingly, when focusing on the multidimensional nature of QOL and how it is affected at the end of life, we should consider the profound influence of social and cultural factors (Qin, 2007). There are a wide range of internationally recognized generic and disease specific quality of life or health status measurement tools. Also there are age group specific instrument of quality of life measurement such as KEEDSCREEN for children and LEIPAD for elderly people. In this article we report the quality of life of elderly people and its relationship with socioeconomic factors.

2. Material and Methods

It was a cross-sectional study carried out in rural area of Sari the capital city of Mezandaran province in northern Iran. LEIPAD questionnaire was used for measuring quality of life of elderly people. A sample of 130 elderly people over 60 years from both sex were interviewed by trained interviewers on their doorstep randomly in 2011. Data collection was continued until we have reached the predetermined sample size. Half of the sample was chosen from mountain area and the second half from plain area. This questionnaire was used in earlier studies among Iranian population (Mohagheghi et al., 2007; Hesamzadeh 2004; Abedi, 1999). LEIPAD consists of 49 items, 31 of which can be grouped into seven 'core instrument scales'. Another 18 items can be grouped into a further five scales, referred to as 'moderator scales'. Each item in the instrument assesses responses along a scale of 0 (best condition) to 3 (worst

condition). Some items of the 'moderator scales' have dichotomy answers so their score is 0 or 1 (55). Collected data were coded and entered in excel program. Inferential statistics and ANOVA were used to analyze the data using SPSS software package.

3. Results

There were 130 respondents of which 58 were male and 62 female. The average age of participants was 71 minimum and maximum of 61 and 90 years respectively. Table 1 show socioeconomic characteristics of respondents.

Table 1. Socioeconomic characteristics of a sample of elderly people living in rural area of Sari Iran 2011

Variables/ frequency		Number	Percent
Sex	Male	58	44.6
	Female	72	55.4
	60-65	36	27.7
Age	66-70	37	28.5
	71-75	26	20.0
	76>	31	23.8
Manuiaga	Couple	90	69.2
Marriage	Single	40	30.8
Source of	Pensioner	48	36.9
income	Self employee	82	63.1
Income	2000000<	49	37.7
category (Iranian	2010000- 4000000	52	40.0
Rials)	4010000>	29	23.3
1*4	Illiterate	121	93.1
literacy	Literate	9	6.9
Location	Mountain	69	53.1
Location	Plain	61	46.9

As table above indicates most of respondents were female, from different age and income categories, living as couple, self employed, illiterate, living in mountain area.

Health status of participants including core instrument scales and moderator scales are presented in table 2. As a matter of cultural norm two questions related to sexual behavior of respondents were ignored reducing the domains of core scales from seven to six items.

Table 2. Health status of a sample of elderly people living in rural area of Sari Iran 2011

	Quality of life domain / Frequency	Mean(St.D)	Percent
	Physical functioning scale 0-15	8.8 (3.2)	58.7
	Self-care scale 0-18	14.6 (3.3)	81.1
Core	Depression and anxiety scale 0-12	7.7 (2.8)	64.2
Instrument	Cognitive functioning scale 0-15	10.0 (3.1)	66.7
Scales	Social functioning scale 0-9	5.3 (1.8)	58.9
	Life satisfaction scale 0-18	11.2 (3.0)	62.2
	Subtotal 0-87	57.6 (12.2)	66.3
	The perceived personality disorder scale 0-6	4.1 (1.5)	68.3
	The anger scale 0-4	3.5 (0.8)	87.5
Moderator Scales	The social desirability scale 0-3	1.9 (0.9)	63.3%
	Self esteem scale 0-3	2.9 (0.4)	96.7
	Trust in God scale 0-3	2.0 (0.2)	66.7
•	Subtotal 0-19	14.4 (2.6)	75.8
_	Total 0-106	72.0 (13.8)	67.9

As table 2 shows the quality of life of respondents in overall was better on core scales (66.3%) than moderator scales (80.0%). Among domains of core scales, physical functioning scale and social functioning scale had better status (about 59%) compared to other domains. Among domains of moderator scales self esteem scale (96.7%) had worst

status where the social desirability (63.3%) and trust in God (66.7%) had the best status. We further analyzed domains of core scales moderator scales and total scales for finding any possible association between those variable with socioeconomic factors indicated in table 1 running one way ANOVA test. The results are presented in table 3.

Table 3. The association between socioeconomic variables and quality of life of elderly people living in rural area of Sari Iran 2011.

Quality of life scales (%) Socioeconomic variables		TCS*	TMS*	TS*
	Female	56.8	14.5	71.3
Sex	Male	58.6	14.1	72.7
	P value	.427	.419	.584
	60-65	63.6	14.9	78.5
	66-70	60.4	14.4	74.8
Age category	71-75	58.2	15.0	73.2
	76>	46.8	13.1	59.9
	P value	.000	.017	.000
	Couple	61.4	14.9	76.3
Marriage	Single	49.0	13.1	62.1
, and the second	P value	.000	.000	.000
	Pensioner	60.2	13.8	74.0
Source of income	Self employee	56.1	14.7	70.7
	P value	.064	.076	.196
Income category (Iranian Rials)	2000000<	54.1	14.4	68.5
	2010000- 4000000	53.9	14.0	67.8
	4010000>	61.8	14.4	76.2
	P value	.001	.754	.005
literacy	Illiterate	56.8	14.4	71.1
	Literate	68.9	13.9	82.8
	P value	.004	.590	.014
	Mountain	56.8	14.8	71.6
Location	Plain	58.3	14.0	72.3
	P value	.468	.072	.764

*Total Core Scales % (TCS); Total Moderator Scales % (TMS) Total Scales % (TS)

As table above indicates among socioeconomic variables studied in this study age, marital status, income and literacy had significantly affected the quality of life of elderly people both in terms of core and moderator scales except for income and literacy that did not change moderator scales significantly. There was no association between sex, source of income, location of residency and quality of life of elderly people neither on core nor on moderator scales.

4. Discussions

In recent decades both the subject of elderly people and quality of life are the focus of many researchers worldwide (Rouhani, 2012; Heydari, 2012). In one side this is because of the number of elderly people particularly in developing countries are

increasing rapidly (Siegel and Doner, 2007; Mortazavi et al., 2012; Jogataee, 2005). For instance in the case of Iran where the number of total population will hardly doubled in 2050, for the same period the number of elderly people has estimated to be six fold (Mehryar, 2004). The other side is that after the successfulness of public health programs and services beside improvement in socioeconomic status of population particularly in developing countries, improvement in health status in terms of morbidity, mortality and length of life has progressed significantly. Putting it into another language, it means that nowadays the gap in terms of health of population is more evident where we take into account the quality of life rather than just the quantity of life.

Health related quality of life studies and quality of life measurement are increasingly carried out universally to indicate the subjective aspects of longevity as the traditional health indicators are failed to cover this important aspects of life (Rouhani, a, 2012; Rouhani, b, 2012). Quality of life measurement is also so important for elderly people as they are at the exposure of many socioeconomic and health conditions that could possibly be a potential risk to jeopardize the quality of their life and satisfaction. In one study in Iran authors find out that supporting elderly people in terms of psychological assistant is a prerequisite for a better quality of life for elderly people (Heydari et al., 2012).

In the near future elderly people from developing countries will account the majority of old age in the world with many of them living in rural area of these countries. Given the inequitable distribution of facilities and social support between urban and rural area of developing countries (Qin, 2007) it should be a big concern for public health policy makers when they think to improve the quality of life of people particularly vulnerable groups such as elderly people. In this study we have found that in the rural area of Iran socioeconomic factors particularly coupling. income, literacy, and age are in particular important in determining the quality of life of elderly people. Correlation between quality of life of elderly people and socio-demographic factors was also found in different studies in Iran (Abbasimoghadam et al., 2009; Tajvar et al., 2008; Vahdaninia et al., 2005; Rafati et al., 2004; Bazrafshan et al., 2008; Herrera Ponce et al., 2008). These evidence need to be translated into appropriate programs to reach these vulnerable groups for adding life to their years (WHO, 2002). The finding of this study is different from another study in the same city that measured quality of life of elderly people in urban area and did not found any significant association between quality of life and socioeconomic status of elderly people. This difference between two studies could be a sign of inequitable distribution of social support between urban and rural area as highlighted by other authors in developing countries (Qin, 2007).

Given the finding of this research that revealed that socioeconomic variable as important factors in determining quality of life of elderly people in rural area and using the available evidence in the literature (Beyaztas et al., 2012; Aghamolaei et al., 2011; Aghanori et al., 2012), it is recommended that appropriate intervention needs to assist vulnerable groups particularly elderly people in rural area to improve the quality of their life.

Acknowledgements:

Authors thank deputy for research and technology at Mazandaran University of medical sciences and its student research committee for the approval and financial support of this study. Also we appreciate the collaboration of all participants who had well cooperation during data collection.

Corresponding Author:

Parsa Zoleikani

Student Research Committee; Health Staff of Sari District Health Centre, Mazandaran University of Medical Sciences, Sari, Iran

E-mail: parsa.z123@yahoo.com

References

- 1. Abbasimoghadam, M.A., Dabiran, S., Safdari, R., Djafarian, K. Quality of life and its relation to sociodemographic factors among elderly people living in Tehran. *Geriatr Gerontol Int.* 2009; 9(3): 270-5.
- 2. Abedi, H. A. A comparison of quality of life of elderly people living at home and residential home in Isfahan. Isfahan University of Medical Sciences. 1999.
- 3. Aghamolaei, T., Sadat Tavafian, S., Zare, S. Determinants of Health Related Quality of Life on People Living in Bandar Abbas, Iran. Iranian J Publ Health, 2011; 40 (2):128-135.
- 4. Aghanori, A., Mahmoodi, M., Salehi, H., Jaeferian, K. Assessing quality of life of elderly people covered by health centers of urban area in Markazy province, Iran in 2011, *Salmand Iranian Journal of Ageing*, 2012; 22 (6): 20-29.
- Banks, J., Marmot, M., Oldfield, Z., & Smith, J. P. (2006). Disease and disadvantage in the United States and in England. Journal of the American Medical Association, 295, 2037-2045.
- Bassuk, S. S., Berkman, L. F., & Amick, B. C. (2002). Socioeconomic status and mortality among the elderly: findings from four US communities. American Journal of Epidemiology, 155, 520-533.
- 7. Bazrafshan, M.R., Hosseini, M.A., Rahgozar, M., Sadat Maddah, S.B. Quality of elderly's life in Shiraz, Jahandidegan club. *Salmand Iranian Journal of Ageing*. 2008; 3(7): 33-41.
- 8. Beyaztas, F. Y., Kurt, G., Bolayir, E. Life satisfaction level of elderly people: a field study in Sivas, Turkey. J Pak Med Assoc. 2012; 62(3): 221-225.
- Herrera Ponce, M. S., Barros Lezaeta, C., Fernández Lorca, M. B. Predictors of Quality of Life in Old Age: A Multivariate Study in

- Chile. Journal of Population Ageing. 2008; 1(1): DOI 10.1007/s12062-011-9043-7.
- 10. Hesamzadeh, A. A comparison of elderly people quality of life living in home and living in elderly people residential home in Tehran, university of rehabilitation, 2004.
- 11. Heydari, J., Rouhani, S., Mohammadpour, R. A. Aging populations' quality of life: An emerging priority for public health system in Iran. Life Science Journal 2012;9(4):1304-1309.
- 12. Huguet, N., Kaplan, S. M., Feeny, D. Socioeconomic status and health-related quality of life among elderly people: Results from the Joint Canada/United States Survey of Health. Social Science & Medicine, 2008; 66: 80-810.
- 13. Jogataee M: The Elderly World Day. *Monthly Magazine of University of Social Welfare and Rehabilitation Science*. 2005; 11: 2.
- 14. Marmot, M. (2004). The status syndrome: How social standing affects our health and longevity. New York, NY: Times book.
- 15. Mehryar, A. H., Ahmad-Nia, S. Age-structural transition in Iran: short and long term consequences of drastic fertility swings during the final decades of twentieth century. CICRED Seminar. Paris, 23-26 February 2004. Available at:http://www.cicred.org/Eng/Seminars/Detail s/Seminars/Popwaves/PopwavesMehryarAhm ad-Nia.pdf
- 16. Mohagheghi, K, S. H., Sajadi, H., Zarea, H., Biglarian, A. A comparison between quality of life of elderly from social security organization and national pension organization living in Qom county 2006, Journal of health management, 2007; 10(27): 49-56.
- Mortazavi, S. S., Mohammad, K., Ardebili, H. E., Beni, R. D., Mahmoodi, M., Keshteli, A. H. Mental disorder prevention and physical activity in Iranian elderly. *Int J Prev Med*. 2012 March; 3(Suppl1): S64–S72. PMCID:PMC339931.
- Mustard, C. A., Derksen, S., Berthelot, J. M., Wolfson, M., & Roos, L. L. (1997). Agespecific education and income gradients in morbidity and mortality in a Canadian province. Social Science & Medicine, 45, 383-397.

- 19. Lynch, J. W., Davey-Smith, G., Kaplan, G. A., & House, J. S. (2000). Income inequality and mortality: importance of individual income, psychosocial environment or material pathways in health. British Medical Journal, 320, 1200-1204.
- Qin, Huali, "Adding Life to Years: Predicting Subjective Quality of Life among Chinese Oldest-Old" (2007). Gerontology Theses. Paper
 - 4. http://digitalarchive.gsu.edu/gerontology_theses/4
- 21. Rafati, N., Mehrabi, Y., Montazeri, A. Quality of life among Kahrizak charity institutionalized elderly people. *Journal of School of Public Health And Institute of Public Health Researches* 2004; 2(3): 67-76.
- 22. Rouhani, S., Kheirkhah, F., Salarieh, I., Abedi, A. Quality of life, its early change and retention in MMT program in Iran: Evidence for policymakers and service providers. Life Science Journal 2012;9(3):2633-2638.
- Rouhani, S., Salarieh, I., Abedi, S., Kheirkhah, F. (2012). Impact of Methadone Maintenance Treatment on the quality of life of opioid dependents in city of Babol. *Journal* of Mazandaran University of Medical Sciences. 22 (87): 47-55.
- Siegel, M., Doner, L. L. Marketing public health: strategies to promote social change, Second Edition. Jones and Bartlett Publications. Boston. USA. 2007.
- 25. Tajvar, M., Arab, M., Montazeri, A. Determinants of health-related quality of life in elderly in Tehran, Iran. *BMC Public Health*. 2008; 8(1): 323.
- Vahdaninia, M.S., Gashtasbi, A., Montazeri, A., Mafton, F. Health quality of life in elderly: population based survey. *Payesh Health Monitor*. 2005; 4: 113-120.
- 27. World Health Organization. *Active Aging: A Policy Framework*. Geneva: World Health Organization, 2002.
- 28. WHOQOL-Group. (1993). Study protocol for the World Health Organization project to develop a Quality of Life assessment instrument (the WHOQOL). *Quality of Life Research*, 2, 153-159.

1/8/2013